

Use only as directed.

- Monthly applications are recommended where there is a high risk of re-infestation, if the pet is allergic to fleas or in situations where tick control is needed.
- For this control to be effective all dogs and cats in the household should be treated simultaneously.
- FRONTLINE SPRAY may be used on pregnant and lactating females and on unweaned puppies/kittens.
- In order to wet the coat down to the skin, it is recommended that a minimum of 3 mL per kg body mass (7.5 mg of active ingredient per kg body mass) to short haired (< 3 cm) animals and 6 mL per kg body mass (15 mg of active ingredient per kg body mass) to long haired ( $\geq$  3 cm) animals be applied.

#### **METHOD OF APPLICATION**

- Bathing or shampooing the animal 2 days before or 2 days after the application is not recommended as this will decrease the duration of activity.
- Mechanical pump spray for external use - each trigger application of the 100 mL bottle gives 0.5 mL and each trigger application of the 250/500 mL bottle gives 1.5 mL spray volume.
- Spray the entire body of the animal, and apply from a distance of approximately 10-20 cm.
- Apply against the lay of the hair and make sure that the entire coat of the animal is wetted
- Ruffle the coat, especially in long haired animals, so that the product penetrates down to the skin.
- Allow to dry naturally.
- Do not towel dry.